

Latin name	Viburnum × carlcephalum		Common name	Fragrant Viburnum	
cultivar	'Chesapeake'		Family	Adoxaceae/Caprifoliaceae	
Year introduced	1981 from 1962 cross		Native range hybrid		
Developed by	Donald Egolf, U.S. National Arboretum				
Patent #/ information					
parentage	Viburnum × carlcephalum 'Cayuga' × Viburnum utile				
Location	Keim East, south			Quantity	1
Source/date	Classic Viburnums 2013		Planting size	Planting date late fall 2013	
Culture					
Sun/shade	full sun to part shade		Soil	acid	
Fertility			Drainage	well-drained	
			Moisture requirements	moist	
Plant characteristics					
Height x Spread	6 × 10		Form	compact, low-spreading	
Flower	slightly fragrant, buds pink				
Color	white		Size	2-3.5"	
Flower Form	snowball		Bloom time	early-mid May	
Leaf color	dark green, semi-evergreen		fall color	red, yellow	
Leaf size			Leaf texture	leathery	
Leaf form			Leaf shape	undulating with a slight twist	
Fruit color	dull red > black		Fruit type	drupe	
Fruit time			Fruit size		
Persistency			Fruit wildlife use		
Bark/stem color					
Bark/stem texture					
Twig					
Bud					
Maintenance					



Pruning	
Wildlife Use	
Native use	
Historical/cultural info	
Folklore/Culture	N/A
Insect Problems	none significant
Disease Problems	none significant
Landscape Uses	
Notes	<p>Not as good as its “sister” cultivar ‘Eskimo’</p> <p>Images: 1 © Viburnum Valley, http://davesgarden.com/guides/pf/showimage/149579/ 2 © Copyright Claire Rosemond 2010 3 Auburn University Horticulture Department 4 © Maurice Laurent 5 © Susan Ladman 6 Forest Farm</p>

