

Latin name	Viburnum			Common name	(hybrid) Viburnum		
cultivar	× 'Emerald Triumph'			Family	Adoxaceae/Caprifoliaceae		
Year introduced	1994			Native range			
Developed by	Harold Pellett, University of Minnesota						
Patent #/ information							
parentage	V. burejaeticum × V. rhytidophylloides 'Alleghany'						
Location	Maxwell Arboretum Viburnum Collection, ID No. 13126				Quantity	3	
Source/date			Planting size	12”-15”	Planting date	9/25/1992	
Culture	adaptable						
Sun/shade	Full sun (best) to part shade		Soil	acid	Drainage	well-drained	
Fertility	Moisture requirements moist						
Plant characteristics							
Height x Spread	8 × 8		Form	rounded			
Flower							
Color	creamy white			Size	2-3”		
Flower Form	slightly rounded cyme			Bloom time	May		
Leaf color	dark green			fall color	yellow, red		
Leaf size	4”		Leaf texture	glossy			
Leaf form	simple		Leaf shape	ovate			
Fruit color	red > black			Fruit type	drupe		
Fruit time	August			Fruit size	3/8-5/8”		
Persistency	Fruit wildlife use						
Bark/stem color							
Bark/stem texture							
Twig							
Bud							
Maintenance							



Pruning	
Wildlife Use	
Native use	
Historical/cultural info	
Folklore/Culture	
Insect Problems	none serious
Disease Problems	highly resistant
Landscape Uses	specimens, groups, masses informal hedges
Notes	<p>See <a href="http://www.hrresearch.org/docs/publications/JEH/JEH_1994/JEH_1994_12_1/JEH%2012-1-59-60.pdf">http://www.hrresearch.org/docs/publications/JEH/JEH_1994/JEH_1994_12_1/JEH%2012-1-59-60.pdf</a></p> <p>Released through the Minnesota Nurserymen's Research Corporation</p> <p>Images: 1, 2, 4, 7 Emily Levine; 3: Grimm's Gardens; 5: © Susan Ladman, Classic Viburnums, 6 Forest Farm</p>

