


Latin name	Viburnum lantana		Common name	Wayfaringtree Viburnum	
cultivar			Family	Adoxaceae/Caprifoliaceae	
Year introduced	Native range Central, southern, western Europe (to Yorkshire in England), NW Africa, SW Asia				
Developed by					
Patent #/ information					
parentage					
Location	Maxwell Arboretum Viburnum Collection, ID No. 317			Quantity	2
Source/date	Planting size		Planting date		
Culture	full sun to partial shade Soil Drainage well-drained Moisture requirements moist, but can tolerate dry soils				
Sun/shade					
Fertility					
Plant characteristics	10'-15' x 10' x 15' Form rounded				
Height x Spread					
Flower	can have unpleasant odor				
Color	white		Size 3" - 5"		
Flower Form	flat-topped cyme		Bloom time May		
Leaf color	dark green to gray-green		fall color	purplish-red	
Leaf size	2"- 5"		Leaf texture	leathery	
Leaf form	simple, uniform serrations		Leaf shape ovate with cordate bases		
Fruit color	red > blue-black		Fruit type	drupe	
Fruit time	August-September		Fruit size	1/3"	
Persistency	Fruit wildlife use numerous birds, Dormouse, Wood Mouse				
Bark/stem color	naked, gray, fuzzy				
Bark/stem texture					
Twig					
Bud					
Maintenance					



Pruning
Wildlife Use
Native use



Historical/cultural info medicinal properties



Folklore/Culture

Insect Problems none serious
Disease Problems none serious

Landscape Uses hedges, borders, or as a deciduous screen

Notes

Images:
1 and 3 Nebraska Statewide Arboretum
2 © Emily Levine
4 Haruta Ovidiu. Copyright © 2014 CC BY-NC 3.0. Bugwood - www.bugwood.org/
5 © Kim Todd
6 USDA-NRCS PLANTS Database / Britton, N.L., and A. Brown. 1913
7 © Copyright Malcolm Storey

