Latin name	Viburnum lentago	Common name Nannyberry V	iburnum	
cultivar		Family <mark>Adoxaceae/Cap</mark>	rifoliaceae	
	1761			
Year introduced	Native range NE and midwestern US; S Canada: N Bruns to SE Sask.			
Developed by				
Patent #/ information				
parentage				
Location	Maxwell Arboretum Viburnum Collection, ID N	Io. 20585 Quantity	1	
Source/date	Plantin			
Culture	adaptable		and the second sec	
Sun/shade	Full Sun to Part Shade, shade tolerant Soil	Drainage well-drained		
Fertility	Moisture	requirements medium, drought tolerant		
	single stemmed tree or large multi-stemmed shrub			
Plant characteristics				
Height x Spread	15-18' × 10-12' Form upright to	arching in old age		
Flower			Cont Contract	
Color	white	Size cymes 3" - 4.5"		
Flower Form	flat-topped cyme	Bloom time mid May		
Leaf color	dark green	fall color yellow, orange, burgundy		
Leaf size	2-4" × 1-2" Leaf text	<sub>ure</sub> glossy		
Leaf form	simply, margin: serrate Leaf sha	аре	5445403	
	pink/rose > bluish black			
Fruit color		Fruit type drupe	To a for the second	
	September - October >	Fruit size .5", oval		
Persistency	very, and abundant Fruit wil	dlife use birds	in the second	
			to de la	
Bark/stem color			The start and st	
Bark/stem texture				
Twig				
Bud				
Maintenance				

Pruning Wildlife Use Native use		NEBRASKA STATEWIJE DOPEČLUM Vitermen Letrage
Historical/cultural info		
Folklore/Culture		
Insect Problems		
Diesease Problems	mildew	N
Landscape Uses	naturalizing, borders, screens	
Notes	moderate to fast growing	
	See: http://plants.usda.gov/factsheet/pdf/fs_vile.pdf	
	1 and 2 © Emily Levine 3 and 7 Rob Routledge, Sault College, Bugwood.org 5 © Nebraska Statewide Arboretum 6	5461657