

Latin name	<i>Viburnum sargentii</i>		Common name	Sargent Viburnum	
cultivar			Family	Adoxaceae/Caprifoliaceae	
Year introduced	1892, Charles S. Sargent		Native range NE Asia		
Developed by					
Patent #/ information					
parentage					
Location	Maxwell Arboretum Viburnum Collection, ID No. 22192				Quantity 1
Source/date			Planting size	Planting date	
Culture	tolerant of wet sites				
Sun/shade	Full Sun to Part Shade		Soil acid	Drainage	
Fertility	Moisture requirements moist				
Plant characteristics					
Height x Spread	10'+ x 10'+		Form	rounded	
Flower	foul-smelling				
Color	creamy white			Size 3-4"	
Flower Form	flat-topped cymes		Bloom time	May	
Leaf color	dark green above, lighter beneath			fall color	yellow/red
Leaf size	2-5" x		Leaf texture	glabrous above, slightly pubesent below	
Leaf form	simple		Leaf shape	tri-lobed with longer middle lobe	
Fruit color	scarlet			Fruit type	drupe
Fruit time	August-October			Fruit size	.3-.5"
Persistency	into winter		Fruit wildlife use		
Bark/stem color	straw to brownish				
Bark/stem texture	glabrous, somewhat corky				
Twig					
Bud	oblong to flask-shaped, .3", scales closely valvate				
Maintenance					



Pruning
Wildlife Use
Native use



Historical/cultural info

Folklore/Culture

Insect Problems
Disease Problems
Landscape Uses

aphids

none serious

groupings, borders

Notes