


# BULBS OF MAXWELL ARBORETUM


botanical name	<i>Anemone blanda</i>		
cultivar	'Blue Shades'		
common name	Grecian Windflower		
family	Ranunculaceae	qty	100+
location	east of creek, north		
origin	Greece, Balkans, Turkey, east to Turkestan		
introduced	1700, 1898	height	3-4"
color	blue		
bloom time	early spring	depth	1-2"
culture	sun>pt. shade; loose sandy soil; tolerant of dry soil		
notes	tuber; soak overnight, plant on edge; Greek: <i>anemos</i> = wind; flat daisy-like flowers, one per stem, fern-likem deeply cut foliage. Dormant after flowering. Will self sow & spread w/o need for division. RHS Award of Garden Merit, 1993		



botanical name	<i>Chionodoxa lucilae</i>		
cultivar	'Pink Giant'		
common name	Glory-of-the-Snow		
family	Hyacinthaceae	qty	
location			
origin	Crete/Cypress/Turkey		
introduced	sp. 1871, cv. 1947	height	6-8"
color	pale pink		
bloom time	early spring	depth	2+"
culture	sun>part shade		
notes	confusion with <i>C. forbesii</i> , formerly Liliaceae; Gr. <i>chion</i> =snow, <i>doxa</i> =glory Star shaped upward facing six petal-like tepals join at the base to form a short tube. Flattened filaments form a cone in the center, making a "white eye." 2-3 narrow, strap-like leaves. (compare to <i>Scilla</i> -tepals not joined)		



botanical name	<i>Crocus ancyrensis</i>	
cultivar	'Golden Bunch'	
common name	Ankara Crocus	
family	Iridaceae	qty 50
location	baldcypress @ bridge	
origin	Turkey	
introduced	species 1879, cv. 1943	height 3"
color	tang.erine yellow	
bloom time	March?	depth 2-3"
culture	sun>pt. shade	
notes	corm Very floriferous; a small very earl cheerful little crocus	



botanical name	<i>Crocus sieberi sublimis</i>	
cultivar	'Tricolor'	
common name	Crocus	
family	Iridaceae	qty 100
location	baldcypress @ bridge	
origin	southern Europe; Peloponnese Mts.	
introduced	1968	height 3"
color	lilac blue w/ white edged golden heart	
bloom time	early spring	depth 2-3"
culture	sun>pt. shade	
notes	RHS Award of Garden Merit, 1993	





botanical nameCrocus vernus

cultivar'Reembrance'

common nameDutch Hybrid Crocus

familyIridaceae

qty

locationalong Loop Road by Gazebo

originEurope

introduced1925

height4-5"


colorpurple

bloom timeearly spring

depth

culture

notescorm; great for forcing



botanical nameErythronium dens-canis

cultivar

common nameDog-tooth Violet

familyLiliaceae

qty13/12

locationshade garden&hosta coll.

originEur., Asia

introduced1596

height6-12"

colorrose-pink

bloom timeMay?

depth3-4"

culturept. shade>shade

noteswoodland soils, mottled foliage



botanical name

cultivar

common name

family  qty

location

origin


introduced  height

color

bloom time  depth

culture

notes



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes






botanical name	<i>Geranium tuberosum</i>	
cultivar		
common name	Tuberous Cranesbill	
family	Geraniaceae	qty 100
location	n-s arb path under hawthorne	
origin	Mediterranean>Iran	
introduced	1596	height 6-8"
color	rose/purple, with darker veins	
bloom time	mid-late sp.	depth 4"
culture	sun>shade	
notes	leaves finely cut, feathery; blooms 1"; Greek <i>geranos</i> =crane; a great little woodland plant, useful for shade	



botanical name	<i>Hyacinthoides hispanica</i> Miller	
cultivar	'Excelsior'	
common name	Spanish Bluebells or Wood Hyacinth	
family	Liliaceae	qty 30
location	mid Vine Arbor area	
origin	SW Eur. (Iberian Peninsula), N. Africa	
introduced	1906	height 10-12"
color	deep blue	
bloom time	late sp.	depth 4"
culture	sun>shade	
notes	slow to establish 15 or more bell-shaped flowers on erect racemes; largest blossoms of all the cultivars	



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes



botanical nameHyacinthus orientalis

cultivar'Sky Jacket'

common nameHyacinth

familyLiliaceae

qty50

locationFleming Flat south

origincentral/western Asia

introduced

height8-10"

colorwisteria blue

bloom timemid spring

depth4-6"

culturesun

notesdrk bl midrib, pale yellow anthers



botanical nameIpheion uniflorum

cultivar'Wisley Blue'

common nameStar Flower

familyLiliaceae

qty50

locationshade bed area, Cornus mas

originSouth America

introduced1961

height3-5"

colorpale blue

bloom timespring

depth2"

culturesun>pt. shade

noteslong bloom period



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes





botanical name

cultivar

common name

family  qty

location

origin

introduced  height 

color

bloom time  depth 

culture

notes 

emergence of fall foliage is a great garden "marker"  
Fragrant bell-shaped flowers on dense racemes;  
short, stout stalks.

botanical name

cultivar

common name

family  qty

location

origin

introduced  height 

color

bloom time  depth

culture

notes 

Div. 9 Poeticus. Award of Merit, Haarlem 1919; 1st  
Class Certificate, Haarlem 1923; RHS: Award of  
Merit 1950; 1st Class Certificate 1968, Award of  
Garden Merit 1993, HC, 1947. Perianth segments  
broadly ovate, spreading, overlapping by 1/2;  
corona shallow, ribbed, mouth wavy, rim crinkled.

botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Div. 3 Small-Cupped Forcing Award, Haarlem 1957, Award of Merit, Haarlem, 1967; RHS FCC, 1977. Perianth segments broadly ovate, broad midrib, overlapping half; corona bowl-shaped, ribbed, mouth expanded, rim tightly frilled.



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Division 3 Small Cup  
Seed parent 'Artillery", pollen parent: 'Avenger'.  
Perianth segments broadly ovate, slightly reflexed, overlapping 1/2; corona shallow, wide-spreading and ribbed with mouth split and overlapping.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Division 6 Cyclamineus. Seed parent: ({'Mitylene' x {N, cyclamineus}} open pollinated. Perianth segments slightly reflexed, margins wavy; corona flared, deeper tone at rim.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Div. 2 Large Cupped. Seed Parent: 'Tryst' Pollen Parent: 'Greenland'. Perianth segments broadly ovate, spreading, smooth, overlapping 1/3; corona lightly ribbed, mouth expanded, wavy. RHS Award of Garden Merit, 1979





botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Div. 6 Cyclamineus Seed parent: 'Golden spur'  
Pollen parent {*N. cyclamineus*}.  
Early Forcing Award, Haarlem: 1928, 1943; RHS  
Award of Merit 1973 & 1st Class Certificate 1974  
(garden); Award of Merit (p) 1977, Award of Garden  
Merit 1993. Petals not as recurved as other Div. 6



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Div. 5 Triandrus/miniature  
RHS Award of Merit, 1938; Award of Garden Merit,  
1995. Seed parent {*N. jonquilla*} Pollen parent {*N. triandrus*} 6-8 flowers per stem; perianth segments  
ovate/oblong, blunt, reflexed, and overlapping at  
base; corona cup-shaped paler than perianth.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Division 6 Cyclamineus; RHS Award of Garden Merit, 1995; Seed parent 'Vulcan' Pollen parent {*N. cyclamineus*}; Perianth segments ovate, dramatically reflexed, overlapping <1/4, corona cylindrical, loosely ribbed, mouth split and overlapping w/ irregularly dentate rim.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height


color

bloom time  depth

culture

notes 

Div. 6 Cyclamineus. Seed parent (('Market Merry' x'Carineer') x'Armada') Pollen parent {*N. cyclamineus*} Perianth segments ovate, strongly reflexed, margins wavy, overlapping 1/3; corona cylindrical, loosely ribbed. RHS Award of Merit 1987, HC(g)1990, Award of Garden Merit, 1995





botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color


bloom time

depth

culture

notes 

Division 7 Jonquilla/miniature. Seed parent 'Ruby'  
Pollen parent {*N. scaberulus*}. 2 flowers per stem;  
very pale overlapping ovate, acute, and spreading  
perianth segments; shallow bowl-shaped ribbed  
corona.



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Division 7 Jonquilla; Seed parent 'Daydream'  
Pollen parent {*N. jonquilla*}. 2-3 flowers per stem,  
perianth segments broadly ovate, overlapping 1/2;  
corona funnel-shaped, loosely ribbed, mouth  
straight or slightly flared, split and overlapping,  
wavy. RHS Award of Garden Merit 1998, Wister  
Award, 2007.





botanical name

cultivar

common name

family

qty

location

origin

introduced

height


color

bloom time

depth

culture

notes



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color


bloom time

depth

culture

notes 

Division 7 Jonquilla (*N.* 'Sweet Charity' x *N. jonquilla*) Fragrant, multiple stems with multiple blooms.



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Div. 6 Cyclamineus?12? Seed parent: 'Cyclataz' self or open polinated. RHS: For cultivation in pots, pans, or bowls: Award of Merit 1956, 1st Class Certificate 1962. Award of Garden Merit 1973, 1st Class Certificate 1974, Award of Garden Merit 1993. 2-3 flowers/stem. Ubiquitous for good reason.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Division 5 Triandrus. 3-4 flowers per stem. Perianth segments ovate/oblong; corona long cup-shaped, ribbed, mouth wavy, rim crenate. Haarlem Award of Merit, 1919, Haarlem Forcing Award 1921.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Division 11 Split Corona. Seed parent: 'April Tears' Pollen parent: 'Baccarat'. 2-3 flowers per stem. Perianth segments broadly ovate, spreading>reflexed, overlapping 1/3-1/2; corona split, rim entire or irregularly crenate. RHS Award of Garden Merit, 2001; Wister Award, 1999





botanical name	<i>Puschkinia scilloides</i> var. <i>libanotica</i> Adams	
cultivar		
common name	Striped Squill, Lebanon Squill	
family	Liliaceae> Hyacinthaceae	qty <input type="text"/>
location	Hosta Collection, near Gazebo	
origin	Caucusus, S Turkey, N Iraq, Iran, Lebanon	
introduced	1805	height 4-6"
color	pale blue	
bloom time	early	depth 2-3"
culture	sun>part shade	
notes	<p>Starry, <i>Scilla</i>-like flowers, pale blue with dark blue midrib stripe. Flowers fused at base to form a tube like <i>Chionodoxa</i>, but the bloom is smaller and never opens flat. Corona is made up of fused filaments creating a small white dome in the center. Leafless stalks. Self-sows.</p>	



botanical name	<i>Scilla bifolia</i> rosea	
cultivar	'Rosea'?	
common name	Two-Leaved Squill	
family	Hyacinthaceae	qty <input type="text"/>
location	Loop Road & Dairy Store Drive, under oak	
origin	S. Europe>Turkey	
introduced		height 3-5"
color	pink	
bloom time	early	depth 2"
culture	sun>shade	
notes	<p>Award of Garden Merit (species), 1993. One-sided racemes of small, delicate, starry flowers. Very pale pink. Strap-like foliage. "rosea" sometimes listed as subspecies, other times as a cultivar.</p>	



botanical name *Scilla litardierei*

cultivar

common name Squill

family Hyacinthaceae      qty

location Loop Road & Dairy Store Drive, under oak

origin S & W Yugoslavia

introduced 1827      height 4-8"

color pale blue violet

bloom time very late      depth

culture

notes RHS Award of garden Merit, 2004.



botanical name *Scilla miczenkoana*

cultivar

common name Squill

family Liliaceae      qty 100

location CYT SW path

origin Asia Minor, N Iran>Caucusus; stoney hillsides

introduced 1931, Tubergen      height 3-4"

color light blue

bloom time early spring      depth 2-3"

culture sun>shade

notes Syn. *S. mischtschenkoana* Light blue with dark blue midrib on exterior; 3-5 flowers per stem, flowers open as they emerge from soil. RHS Award of Garden Merit, 2004. Confusion surrounds the species nomenclature.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height 

color

bloom time  depth 

culture

notes 

naturalizes. Strap-like leaves emerge after flowering; pendent, bell-shaped, striking blue flowers are born on loose one-sided racemes. RHS Award of Garden Merit, 1993

A photograph showing a dense cluster of small, bell-shaped blue flowers of Scilla siberica. The flowers are hanging from green, strap-like leaves. The background is dark and out of focus. A red border surrounds the image, and a pink watermark "@Kim Todd" is visible in the lower-left corner.

botanical name

cultivar

common name

family  qty

location

origin

introduced  height 

color

bloom time  depth 

culture

notes 

Award of Garden Merit (species) 1993. 4-6 narrow, strap-like leaves emerge after flowering. One sided racemes bearing 2-5 nodding, bell-shaped flowers. Intense deep blue.

A close-up photograph of a single, larger blue flower of Scilla siberica. The flower has six distinct petals and a green center with visible stamens. The background is blurred, showing green foliage. A red border surrounds the image.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height


color

bloom time  depth

culture

notes 

Div. XI Kaufmanniana Bloom resembles a waterlily.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height


color

bloom time  depth

culture

notes 

Div. I Single Early. Fragrant



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Div. IV Darwin Hybrid. Reliably perennial; classic tulip form and color



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes 

Div. XI Kaufmanniana. Mottled purple foliage; tall, thin flower form.



botanical name

cultivar

common name

family

qty

location

origin

introduced

height

color

bloom time

depth

culture

notes



botanical name

cultivar

common name

family

qty

location

origin

introduced

height


color

bloom time

depth

culture

notes





botanical name

cultivar

common name

family  qty

location

origin

introduced  height 

color

bloom time  depth 

culture

notes 

Div. V Single Late

botanical name

cultivar

common name

family  qty

location

origin

introduced  height 

color

bloom time  depth

culture

notes 

Darwin Hybrid. Long-lasting

botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Stoloniferous. Leaves form a rosette at base of plant; starry yellow flowers with white tips.



botanical name

cultivar

common name

family  qty

location

origin

introduced  height

color

bloom time  depth

culture

notes 

Div. XIV Miscellaneous  
An exceptionally cheery little tulip.

